

13
LEVEL II



AD A108900

STUDIES ON THE MECHANISM OF ACTION OF THE IN VITRO PGB_x EFFECT
III. THE INTERACTION BETWEEN PGB_x AND RAT LIVER MITOCHONDRIA

Herman W. Shmukler, Ph.D.
Sara F. Kwong, M.S.
Biochemistry Research Team
Aircraft and Crew Systems Technology Directorate
Naval Air Development Center
Warminster, Pennsylvania 18974

Margaret G. Zawryt, B.S.
Elaine F. Soffer, B.S.
Hahnemann Medical College
Philadelphia, Pennsylvania 19102

17 September 1981

6217 N
Phase Report
Airtask No. F58527803
Work Unit No. EH810

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

Prepared for
Office of Naval Research
Department of the Navy
Arlington, Virginia 22217

DTIC
ELECTE
S DEC 29 1981 D
D

2
398532
8112 28047

DTIC FILE COPY


NOTICES

REPORT NUMBERING SYSTEM – The numbering of technical project reports issued by the Naval Air Development Center is arranged for specific identification purposes. Each number consists of the Center acronym, the calendar year in which the number was assigned, the sequence number of the report within the specific calendar year, and the official 2-digit correspondence code of the Command Office or the Functional Directorate responsible for the report. For example: Report No. NADC-78015-20 indicates the fifteenth Center report for the year 1978, and prepared by the Systems Directorate. The numerical codes are as follows:

CODE	OFFICE OR DIRECTORATE
00	Commander, Naval Air Development Center
01	Technical Director, Naval Air Development Center
02	Comptroller
10	Directorate Command Projects
20	Systems Directorate
30	Sensors & Avionics Technology Directorate
40	Communication & Navigation Technology Directorate
50	Software Computer Directorate
60	Aircraft & Crew Systems Technology Directorate
70	Planning Assessment Resources
80	Engineering Support Group

PRODUCT ENDORSEMENT – The discussion or instructions concerning commercial products herein do not constitute an endorsement by the Government nor do they convey or imply the license or right to use such products.

APPROVED BY:


J. R. WOODS
CDR USN

DATE:

11/25/81

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NADC-81243-60	2. GOVT ACCESSION NO. AD-A108900	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Studies on the Mechanism of Action of the <u>in vitro</u> PGB _x Effect III. The Interaction Between PGB _x and Rat Liver Mitochondria		5. TYPE OF REPORT & PERIOD COVERED Phase Report
7. AUTHOR(s) Herman W. Shmukler, Sara F. Kwong, Margaret G. Zawryt and Elaine F. Soffer		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Air Development Center Aircraft and Crew Systems Technology Directorate Warminster, PA 18974		8. CONTRACT OR GRANT NUMBER(s) N00014-81-WR10070
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Airtask: ^W F58527803 Work Unit No.: EH810
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 17 September 1981
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) PGB _x Rat liver mitochondria Oxidative phosphorylation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Tritium labelled PGB _x -active oligomers were synthesized by alkaline polymerization of ³ H-PGB ₁ . When added to PGB _x assay system non-stoichiometric amount of ³ H-PGB ₁ were bound to RLM. The degree of binding was dependent upon the concentration of ³ H-PGB ₁ in the reaction. On the assumption that ³ H-PGB ₁ and PGB _x are similar or identical in structure, then no correlation between PGB _x effect and PGB _x binding by RLM could be found.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

NADC-81243-60

TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION	1
METHODS AND MATERIALS	1
EXPERIMENTAL	2
DISCUSSION	4
FIGURE 1	5
FIGURE 2	6
REFERENCES	7

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
7	

DTIC
ELECTE
S **D**
DEC 29 1981
D

INTRODUCTION

The therapeutic use of PGB_x (1, 2, 3) in the treatment of human ischemic pathologies was suggested by successful animal experiments in a number of laboratories (4, 5, 6, 7, 8, 9, 10). Before human trials maybe attempted, it is necessary to first know the mechanism of the in vivo action of PGB_x . Towards this end studies have been underway in this laboratory to elucidate the in vitro mechanism of the PGB_x effect on phosphorylation activity of degraded RLM, in the hope that this would then lead to an understanding of the in vivo mechanisms of action.

In an earlier report (11) it was shown that the sequence in which PGB_x was added to the in vitro PGB_x assay system was important. When PGB_x was added to already degraded RLM, no maintenance of high phosphorylation activity was observed. Only when RLM were degraded in the presence of PGB_x could the PGB_x effect be demonstrated. These results suggested that RLM reacts with PGB_x to form a complex that protects the RLM from exposure to the hypotonic degradation medium. In this paper more direct evidence for this interaction is reported.

METHODS AND MATERIALS

PGB_1 -methyl ester, 15-keto PGB_1 -methyl ester and PGB_x Type II were synthesized as described previously (2, 3). ^3H - PGB_1 , labelled in the 5, 6 position, was purchased from New England Nuclear Corporation (Boston, MA). Radioactivity was measured with a Beckman Model LS-200B Liquid Scintillation Syst (Fullerton, CA) using "Bray's Mixture" (12) (0.2g dimethyl POPOP, 4g PPO, 60g naphthalene, 100 ml methanol, 20 ml ethylene glycol, and dioxane to make one liter). RLM phosphorylation activity was measured as previously reported (2, 3).

Note: Abbreviations used in this report are: RLM, rat liver mitochondria; P_i , inorganic phosphate; PG-, prostaglandin; PGB_x , oligomeric mixture from alkaline polymerization of 15-keto PGB_1 ; PGB_{1x} , oligomeric mixture from alkaline polymerization of PGB_1 ; PPO, diphenyloxazole; dimethyl POPOP, 1,4-BIS (2-(4-methyl-5-phenyloxazoly 1)) benzene

EXPERIMENTAL

The best method for the measurement of the PGB_x -RLM complex entails the separation of the RLM from the remainder of the reaction medium and then the analysis of the RLM for its PGB_x content. The separation of the RLM is easily accomplished by centrifugation; the analysis of the PGB_x content however is not easily accomplished. At the present time, methods for the quantitative analysis for PGB_x is based on its characteristic UV absorption or fluorescence spectra. However, these optical measurements cannot be used with RLM directly because of their insolubility. An alternative procedure then would be to analyze the clear supernatant solutions from reaction media in which RLM was first sedimented. Any decrease in the soluble PGB_x content from that amount of PGB_x added would then be a measure of the PGB_x bound to RLM. Attempts to use the optical methods for PGB_x analysis of the clear supernatant solutions were not successful, because the constituents of the assay medium also exhibited similar UV absorption and fluorescence spectra, and thus interfered with the measurement of trace amounts of PGB_x .

It appeared obvious then, that the successful determination of PGB_x bound to RLM would require a very sensitive and specific analytical method. It is well known that the use of radioactive tracers permits the specific analysis of trace amounts of material, and this method then appeared to be the method of choice for this study.

Synthesis of $^3\text{H-PGB}_{1x}$: The synthesis of radioactive labelled PGB_x from 15-keto PGB_1 was not feasible because of commercial unavailability. Since $^3\text{H-PGB}_1$ is available commercially, and since PGB_1 had been converted to active PGB_x by Polis et al (1) it was decided to prepare an oligomer of $^3\text{H-PGB}_1$ that would exhibit in vitro PGB_x properties. This oligomeric mixture was designated $^3\text{H-PGB}_{1x}$ to show the identity of the precursor used. $^3\text{H-PGB}_{1x}$ was synthesized by dissolving a mixture of 50 μC of $^3\text{H-PGB}_1$ and 10 μg of PGB_1 -methyl ester in 0.4 ml of ethanolic-KOH (50% ethanol - 1.0N KOH) and heating the mixture in a stoppered flask at 50° for five hours. The polymerization reaction was stopped by adding 10 ml of H_2O and acidifying pH 3.0 with HCl. The PGB_x synthetic conditions used in this study were found to be optimal from preliminary test polymerizations of unlabelled PGB_1 -methyl ester in which the time and temperature of the reaction were varied. $^3\text{H-PGB}_{1x}$ was purified by extraction into ethyl acetate (2 extractions, 20 ml each), washing the combined ethyl acetate layers with H_2O (4 times with, 15 ml) to remove excess acid, and centrifugation to remove insoluble material. The solvent was removed by flash-evaporation at 50° and the residue dried to constant weight. Yield: 9.1mg (91%); specific


activity, 3194cpm/ μ g. For use in this study, the $^3\text{H-PGB}_{1x}$ was dissolved in 25% ethanol-0.0025M NaHCO_3 to a final concentration of 10mg/ml and stored at 0° until used.

$^3\text{H-PGB}_{1x}$ was assayed for the in vitro PGB_x effect as described previously (2, 3) over a concentration range of 1.36 to 36.36 μ g per ml of reaction medium. These results and the results of the assay of PGB_x Type II are shown in Figure 1. The PGB_x activity of $^3\text{H-PGB}_{1x}$ was almost equivalent to that of PGB_x Type II except that at the higher concentration, $^3\text{H-PGB}_{1x}$ exhibited less inhibitor effect.

Binding of $^3\text{H-PGB}_{1x}$ by RLM: To four beakers containing the reagents of the degradation mixture of the PGB_x assay system (2, 3), aliquots of stock $^3\text{H-PGB}_{1x}$ were added so that each beaker contained either 1 to 15 μ g/ml reaction. After equilibration at 27° , 4mg of RLM, aged four days at 0° in 0.3M sucrose, were added and the mixture incubated at 27° for eight minutes. At the end of this time period, the mixture was centrifuged at 8000g for ten minutes to sediment the RLM. The radioactivity in the supernatant was determined by adding 0.1 ml of clear supernatant to 10 ml of Bray's scintillation cocktail and counting the radioactivity for at least two minutes. In another identical series of tests, the nucleotide mixture of stage 2 of the PGB_x assay system was added and the mixture incubated for 20 minutes. At the end of this time the mixture was centrifuged at 8000g for ten minutes and aliquots were removed for radioactivity counting and in addition, another aliquot was deproteinized and the phosphorylation activity determined as described previously (2, 3). The phosphorylation results are plotted in Figure 1. The amount of $^3\text{H-PGB}_{1x}$ in the soluble fraction of each reaction mixture was calculated using the specific activity of $^3\text{H-PGB}_{1x}$. By subtracting the soluble $^3\text{H-PGB}_{1x}$ from the total $^3\text{H-PGB}_{1x}$ added, the amount of $^3\text{H-PGB}_{1x}$ sedimenting with the RLM was calculated. Figure 2 shows the plot of $^3\text{H-PGB}_{1x}$ bound to RLM in terms of $\mu\text{g}/\mu\text{g}$ RLM as a function of total $^3\text{H-PGB}_{1x}$ added. The solid line represents the degradation stage only, while the dashed line represents the total PGB_x assay system. Although the phosphorylation reactions appear to cause a discharge of bound $^3\text{H-PGB}_{1x}$ as evidenced by the lower amounts of $^3\text{H-PGB}_{1x}$ bound, the overall picture actually suggests no change in bound $^3\text{H-PGB}_{1x}$ and the lowered results maybe experimental error. The amount of $^3\text{H-PGB}_{1x}$ bound to RLM appears to be dependent upon the amount of $^3\text{H-PGB}_{1x}$ added to the system. The maximum PGB_x effect takes place at the level between 10-20 μ g $^3\text{H-PGB}_{1x}$. At higher levels, where $^3\text{H-PGB}_{1x}$ decreases phosphorylation, a greater amount is bound to the RLM.

DISCUSSION

On the assumption that the PGB_x -active oligomers formed during the alkaline polymerization of PGB_1 (PGB_{1x}) are similar or identical to PGB_x oligomers, then the binding of PGB_{1x} by RLM also means the RLM binds PGB_x in a similar fashion. On this basis then, RLM binds non-stoichiometric amounts of PGB_x and that the degree of binding is dependent upon the concentration of PGB_x added. As to whether the RLM- PGB_x complex is important in the PGB_x effect on RLM oxidative phosphorylation, is not answered by the results of this study. The results, however, do show that a correlation does not exist between the PGB_x effect and the degree of PGB_x binding to RLM, since at low PGB_x : RLM proportions, the phosphorylating activity is high; at high PGB_x : RLM proportions, the phosphorylating activity is low.






FIGURE 2: the binding of $^3\text{H-PCB1x}$ to KLM during the assay of phosphorylation activity. Samples of $^3\text{H-PCB1x}$ synthesized as described under "pg. look" were tested for PCB_x activity as described previously (2, 3). At the end of the degradation step (8 minutes, at 27°) the reaction mixture was centrifuged at 0° for 10 minutes at 8000g. The radioactivity in the supernatant layer was measured as described under "pg. look". The $^3\text{H-PCB1x}$ bound to KLM was calculated by subtracting the soluble radioactivity from the added radioactivity, and converting to μg by applying the specific activity factor. The solid line curve is the $^3\text{H-PCB1x}$ bound to KLM as a function of added $^3\text{H-PCB1x}$ during the degradation stage of the PCB_x assay. The dashed line curve is the $^3\text{H-PCB1x}$ bound to KLM as a function of added $^3\text{H-PCB1x}$ after stages I and II of the PCB_x assay (2, 3).

REFERENCES

1. Polis, B. D., A. M. Grandizio, E. Polis: Some in vitro and in vivo Effects of a New Prostaglandin Derivative. Neurohumoral and Metabolic Aspects of Injury. Adv. in Exper. Med. and Biol., Plenum Press, New York 33:213-220 (1973).
2. Polis, B. D., S. F. Kwong, E. Polis and G. Nelson: Studies on PGB_x: A Polymeric Derivative of Prostaglandin B₁: I - Synthesis and Purification of PGB_x. Report No. NADC-78235-60 (1978).
3. Polis, B. D., E. Polis and S. F. Kwong: Protection and Reactivation of Oxidative-Phosphorylation in Mitochondria by a Stable Free-Radical Prostaglandin Polymer (PGB_x). Proc. Natl. Acad. Sci., 76:1598-1602 (1979).
4. Angelakos, E. T., B. D. Polis and R. L. Riley: Recovery After Coronary Ligation and Fibrillation in Primates Treated with a Prostaglandin Derivative. 47th Scientific Sessions American Heart Association, Abstracts, Dallas Texas (1974).
5. Angelakos, E. T., B. D. Polis and R. L. Riley: Protection by a Prostaglandin Derivative from Mortality after Coronary Ligation from Ventricular Fibrillation in Primates. 6th International Congress of Pharmacology Abstracts, Helsinki, Finland (1975).
6. Angelakos, E. T., R. L. Riley and B. D. Polis: Recovery of Monkeys from Cardiogenic Shock After Myocardial Infarction with Ventricular Fibrillation. Effects of PGB_x. Report No. NADC-77308-60 (1977).
7. Kolata, R. J.: The Effect of PGB_x on Neurological Recovery from Cerebral Ischemia in Rabbits. Masters Thesis, Univ. of Penna. Veterinary School (1977).
8. Kolata, R. J. and B. D. Polis: Facilitation of Recovery from Ischemic Brain Damage in Rabbits by Polymeric Prostaglandin PGB_x, a Mitochondrial Protective Agent. Physiol. Chem. and Physics 12, 551 (1980).
9. Yamazaki, H., M. M. Bodenheimer, V. S. Banka, J. Lewandowski and R. H. Helfant: The Effect of a New Prostaglandin (PGB_x) on Length-tension Relationship Following Partial Coronary Occlusion and Reperfusion. Amer. Heart. Assn., Abstracts 51st Scientific Session, Dallas, Texas, (1978).

NADC-81243-60

10. Moss, G., T. Magliochetti and A. Quarmby: Immediate Restoration of Central Nervous System Autonomic Cardio-pulmonary Control: Survival of "Lethal" Cerebral Hypoxia by Treatment with PGB_x. Surgical Forum, 21:513, (1978).
11. Shmukler, H. W., M. G. Zawryt and E. Soffer: Report No. NADC-81223-60 (1981).
12. Instruction Manual LS-200B (Page 2-10). Beckman Instruments, Fullerton, CA.

DISTRIBUTION LIST

REPORT NO. NADC-81243-60

CONTRACT NO. N00014-81-WR1007-0

No. of Copies

Defense Technical Information Center, Building 5, Cameron Station, Alexandria, VA 22314	13
Naval Research Laboratory, Attn: Technical Information Division (Code 2627), Washington, DC 20375	6
Office of Naval Research, Attn: Code 102IP (ONRL DOC), 800 N. Quincy Street, Arlington, VA 22217	6
Office of Naval Research, Biophysics Program (Code 444), 800 N. Quincy Street, Arlington, VA 22217	3
Office of Naval Research (Code 200), 800 N. Quincy Street, Arlington, VA 22217	1
Office of Naval Research, Branch Office, Building 114, Section D, 600 Summer Street, Boston, MA 02210	2
Office of Naval Research, Branch Office, 536 S. Clark Street, Chicago, IL 60605	1
Office of Naval Research, Branch Office, 1030 E. Greet Street, Pasadena, CA 91106	1
Naval Medical Research & Development Command, National Naval Medical Center, Bethesda, MD 20014	1
Chief, Bureau of Medicine & Surgery, Department of the Navy, Washington, DC 20375	1
Naval Medical Research Institute, Tech. Ref. Library, National Naval Medical Center, Bethesda, MD 20014	2
Naval Medical Research Unit No. 2, Box 14, APO, San Francisco, CA 96263	1
Naval Medical Research Unit No. 3, FPO, NY 09527	1
Submarine Medical Research Laboratory, Naval Submarine Base, New London, Groton, CT 06342	1
Naval Aerospace Medical Research Institute, Scientific Library, Naval Aerospace Medical Center, Pensacola, FL 32512	1
Naval Biosciences Laboratory, Building 844, Naval Supply Center, Oakland, CA 94625	1
Army Research Office, P.O. Box 12211, Research Triangle Park, NC 27709	1
Air Force Office of Scientific Research, Directorate of Life Sciences, Bolling Air Force Base, Washington, DC 20332	1
Army Medical Research & Development Command, Forrestal Building, Washington, DC 20314	1
Department of the Army, U.S. Army Science & Technology Center, Far East, APO, San Francisco, CA 06328	1